

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mouse anti Carcinoembryonic Antigen / CEA, IgG1, Clone: [PARLAM 4], Monoclonal NMB-MUB0332P**

|                          |   |
|--------------------------|---|
| Artikelname              | Mouse anti Carcinoembryonic Antigen / CEA, IgG1, Clone: [PARLAM 4], Monoclonal  |
| Artikelnummer            | NMB-MUB0332P  |
| Hersteller Artikelnummer | MUB0332P  |
| Alternativnummer         | NMB-MUB0332P  |
| Hersteller               | NordicMubio   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, ICC, IHC-Fr, IHC-P, WB  |
| Spezies Reaktivität      | Human   |
| Produktbeschreibung      | This monoclonal antibody is reactive with Human carcinoembryonic antigen (CEA), a tumour associated antigen with oncofetal characteristics. Although CEA can be found in tissues of non-neoplastic diseases and normal epithelia, it occurs also in a larg... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PARLAM 4]  |
| Isotyp                   | IgG1  |
| UniProt                  | <a href="#">P06731</a>  |

|                        |   |
|------------------------|---|
| Puffer                 | Each vial contains 100 ul 1 mg/ml purified antibody in PBS containing 0.09% sodium azide.   |
| Quelle                 | PARLAM 4 is a mouse monoclonal IgG1 antibody derived by fusion of Sp 2/0 Ag 14 Mouse myeloma cells with spleen cells from a BABL/c Mouse immunized with isolated Human CEA. The immunogen has been isolated from Human colonic carcinoma cells.   |
| Anwendungsbeschreibung | PARLAM 4 is useful for flow cytometry, immunoblotting, immunocytochemistry on methanol fixed cells and immunohistochemistry on frozen tissues when using a PBS buffer containing 0.1 mM CaCl <sub>2</sub> and 0.1 mM MgCl <sub>2</sub> . The antibody is also reactive in formalin-fixed and paraffin-embedded tissue sections after treatment with citrate buffer pH 6.0 in an autoclave. Human colon carcinoma tissue is used as positive control. Optimal antibody dilution should be determined by titration, we recommend a 1:25 - 1:100 dilution for immunohistochemistry with avidin-biotinylated Horseradish peroxidase complex (ABC) as detection reagent, and 1:100 - 1:500 for immunoblotting application. |