

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Neurofilament 200 kD, IgM, Clone: [RNF402], Monoclonal NMB-MUB1307P

Artikelname	Mouse anti Neurofilament 200 kD, IgM, Clone: [RNF402], Monoclonal
Artikelnummer	NMB-MUB1307P
Hersteller Artikelnummer	MUB1307P
Alternativnummer	NMB-MUB1307P
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	ICC, IHC-Fr, IHC-P, WB
Spezies Reaktivität	Bovine, Canine, Gallus, Guinea pig, Hamster, Human, Monkey, Mouse, Rabbit, Rat, Sheep, Xenopus
Produktbeschreibung	Like most other intermediate filament proteins (IFPs), the expression of the different neuronal IFPs is both tissue-specific and developmentally regulated. The neurofilament (NF) triplet proteins (70, 160, and 200 kDa) occur in both the central and p...
Klonalität	Monoclonal
Konzentration	1 mg/ml
Klon-Bezeichnung	[RNF402]
Isotyp	IgM
UniProt	P12036

Puffer	Each vial contains 0.1 ml monoclonal antibody in PBS containing 0.09% sodium azide. The antibody is purified by ammonium sulphate precipitation and will therefore contain fetal calf serum derived protein contaminants.
Quelle	RNF402 is a Mouse monoclonal IgM antibody derived by fusion of SP2/0-Ag14 Mouse myeloma cells with spleen cells from a BALB/c Mouse immunized with a neurofilament preparation of calf brain tissue.
Reinheit	Ammonium sulphate precipitation.
Formel	Each vial contains 0.1 ml monoclonal antibody in PBS containing 0.09% sodium azide. The antibody is purified by ammonium sulphate precipitation and will therefore contain fetal calf serum derived protein contaminants
Anwendungsbeschreibung	RNF402 is suitable for immunoblotting, immunocytochemistry on permeabilised cells and immunohistochemistry on frozen and paraffin-embedded tissues. Optimal antibody dilution should be determined by titration, recommended range is 1:10 - 1:100 for immunohistochemistry with avidin-biotinylated Horseradish peroxidase complex (ABC) as detection reagent, and 1:25 - 1:250 for immunoblotting applications.