

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse IgG1 Negative Control, Clone: [ZX3], Unconjugated, Monoclonal NMB-0G11

Artikelname	Mouse IgG1 Negative Control, Clone: [ZX3], Unconjugated, Monoclonal
Artikelnummer	NMB-0G11
Hersteller Artikelnummer	0G11
Alternativnummer	NMB-0G11
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC
Konjugation	Unconjugated
Produktbeschreibung	This monoclonal antibody to IgG1 is a negative control for Immunofluorescence staining with Pure, FITC, Biotin, PE and APC mouse monoclonal antibodies of the IgG1 subclass. It is used to monitor the level of non-specific binding of mouse monoclonal a...
Klonalität	Monoclonal
Konzentration	See vial for concentration
Klon-Bezeichnung	[ZX3]
Isotyp	IgG1
UniProt	P07750

Puffer	Provided as solution in phosphate buffered saline with 0.08% sodium azide
Reinheit	Protein A/G Chromatography
Formulierung	Unconjugated
Anwendungsbeschreibung	Consult the appropriate fact sheet to determine the amount of antibodies to be used as a control for PBMC or Whole Blood. PBMC: Add 10 µl of MAB/10 6 PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of Whole Blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope. ALLOPHYCOCYANIN: (APC) conjugates are analyzed in multi-color flow cytometry with instruments equipped with a second laser and proper filters. Laser excitation is at 633 nm with a Helium Neon (HeNe) laser or a 600-640 nm (633nm) range for a Dye laser. Peak fluorescence emission is at 660 nm. RPE-Cy-5 +: Excites at 488nm and emits at 670nm. Store protected from light. Optimal concentration should be evaluated by serial dilutions.