

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat anti Chicken IgG IgA IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GACH/IG

Artikelname	Goat anti Chicken IgG IgA IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GACH/IG
Hersteller Artikelnummer	GaCh/Ig
Alternativnummer	NMB-GACH/IG
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Gallus
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins. It does not react with any non-Ig protein in chicken serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunoelectropho...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat seru
Quelle	Purified normal Ig fractions isolated from pooled chicken serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.