

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Dog IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GAD/IGG(H+L)/7S

| | |
|----------------------------------|---|
| Artikelname | Goat anti Dog IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAD/IGG(H+L)/7S |
| Hersteller Artikelnummer | GAD/IgG(H+L)/7S |
| Alternativnummer | NMB-GAD/IGG(H+L)/7S |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Spezies Reaktivität | Canine |
| Konjugation | Unconjugated |
| Format | IgG |
| Spezifität | IgG (H+L) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in dog serum, as t... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |

| | |
|------------------------|---|
| Puffer | Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added. It is reconstituted by adding 1 ml sterile distilled water |
| Quelle | Purified normal IgG isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added. |
| Antibody Type | Secondary Antibody |
| Anwendungsbeschreibung | Indirect immunofluorescence, ELISA, Dot blot, Immunoblotting. |