

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Horse IgG (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAHO/IGG(FC)

Artikelname	Goat anti Horse IgG (Fc specific), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAHO/IGG(FC)
Hersteller Artikelnummer	GAHo/IgG(Fc)
Alternativnummer	NMB-GAHO/IGG(FC)
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Applikation	IP
Spezies Reaktivität	Equine
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal horse serum a single precipitin line is obtained which shows ...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a
Quelle	Highly purified normal IgG isolated from pooled horse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat and serially diluted in 2 mm diameter peripheral wells).