

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human alpha-2 macroglobulin, Clone: [Polyclonal], Monoclonal NMB-GAHU/A2M

Artikelname	Goat anti Human alpha-2 macroglobulin, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAHU/A2M
Hersteller Artikelnummer	GAHu/A2M
Alternativnummer	NMB-GAHU/A2M
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Produktbeschreibung	The reactivity of the antiserum is restricted to human alpha-2 macroglobulin. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal human plasma a single precipitin line is obtained which shows a r...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
UniProt	P01023
Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added as it may interfere with the antibody activity. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum.

Quelle	Alpha-2 macroglobulin(clotting factor I) is the major protein in the alpha-2 region of human serum after separation by immunoelectrophoresis. The concentration in normal plasma is about 2.5 mg/ml. It is a large polymer and a strong antigen. It forms about
Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added as it may interfere with the antibody activity. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum.
Anwendungsbeschreibung	Precipitation assays. The lyophilized antiserum is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4C, not refrozen, and preferably used the same day.