

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti human Factor VIII related antigen (VWD), Clone: [Polyclonal], Monoclonal NMB-GAHU/FVIII VWD

Artikelname	Goat anti human Factor VIII related antigen (VWD), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAHU/FVIII VWD
Hersteller Artikelnummer	GAHu/FVIII VWD
Alternativnummer	NMB-GAHU/FVIII VWD
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Produktbeschreibung	The defined antibody reactivity is restricted to VWF. In immunoelectrophoresis, bi-dimensional electrophoresis and radial immunodiffusion (Ouchterlony) against normal plasma, a single precipitin line is obtained which shows a reaction of identity wit...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
Puffer	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. IgG protein concentration in the antiserum is 10 mg/ml. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water.

Quelle	Human FVIII procoagulant activity (FVIIIIC) is carried by a polypeptide non-covalently bound to a large carrier molecule (unit molecular weight 250 kD) known as the Von Willebrand factor (VWF). FVIII/VWF exists in plasma as a series of polymers with molecu
Formel	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. IgG protein concentration in the antiserum is 10 mg/ml. No foreign proteins added.
Anwendungsbeschreibung	The antiserum concentration required in the gel is normally between 1 and 2%. In immunologic determinations of FVIII/VWF, plasma samples and all assay components must contain EDTA to stabilize the proteins The antiserum is standardized for use in the electroimmunodiffusion (EID, Laurell) test procedure for the quantitative determination of FVIII/VWF as scribed in the Nordic Recommended Working Procedure.