

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human IgE (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAHU/IGE(FC)

| | |
|----------------------------------|---|
| Artikelname | Goat anti Human IgE (Fc specific), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAHU/IGE(FC) |
| Hersteller Artikelnummer | GAHu/IgE(Fc) |
| Alternativnummer | NMB-GAHU/IGE(FC) |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Konjugation | Unconjugated |
| Format | Antiserum |
| Spezifität | IgE |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | The reactivity of the antiserum is restricted to the Fc part of the IgE molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal human serum containing monoclonal IgE a single precipitin lin... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |

| | |
|------------------------|--|
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized ant |
| Quelle | Highly purified monoclonal IgE isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Antibody Type | Secondary Antibody |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat and serially diluted in 2 mm diameter peripheral wells). |