

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human plasminogen, Clone: [Polyclonal], Monoclonal NMB-GAHU/PLG

| | |
|--------------------------|---|
| Artikelname | Goat anti Human plasminogen, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAHU/PLG |
| Hersteller Artikelnummer | GAHu/Plg |
| Alternativnummer | NMB-GAHU/PLG |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Produktbeschreibung | The defined antibody reactivity is restricted to plasminogen as tested at the level of sensitivity of immuno-precipitation techniques. A single precipitin line is obtained in bidimensional electrophoresis, immunoelectrophoresis and double radial immu... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| UniProt | P00747 |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized anti |
| Quelle | Native plasminogen is a single polypeptide chain, synthesized in the liver. Its molecular weight has been reported to be 81,000 and 92,000. Part of the molecule contains the active serine esterase of plasmin. On activation it is converted to two polypepti |

| | |
|------------------------|--|
| Formel | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. In electroimmunodiffusion the amount of antiserum required in the agarose gel is usually between 1 and 2% depending on the test arrangement. |