

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human secretory component (free and bound) , conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-GAHU/SC/FITC

Artikelname	Goat anti Human secretory component (free and bound) , conjugated with FITC, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAHU/SC/FITC
Hersteller Artikelnummer	GAHu/SC/FITC
Alternativnummer	NMB-GAHU/SC/FITC
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Human
Konjugation	FITC
Format	IgG
Spezifität	Secretory Component - free and bound determinants
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Tested in immunoelectrophoresis, double radial immunodiffusion and ELISA against a panel of appropriate secretions and purified Ig isotypes. The antiserum reacts with both bound secretory component (secretory IgA) and with the free SC present in huma...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Fluorochrome-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.
Quelle	Secretory component is present in human secretions bound to secretory IgA (sIgA) and in free form. Secretory IgA (sIgA) functions as a dimer or polymer and accounts for almost all specific mucosal antibody activity. A molecule of sIgA is made up of two mo
Formulierung	FITC
Formel	Fluorochrome-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. The lyophilized conjugate is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided.