

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat anti Mouse Ig lambda light chain (free and bound), Clone: [Polyclonal], Monoclonal NMB-GAM/BJL(SD+HD)

|                                  |   |
|----------------------------------|---|
| Artikelname                      | Goat anti Mouse Ig lambda light chain (free and bound), Clone:<br>[Polyclonal], Monoclonal  |
| Artikelnummer                    | NMB-GAM/BJL(SD+HD)  |
| Hersteller Artikelnummer         | GAM/BJL(SD+HD)  |
| Alternativnummer                 | NMB-GAM/BJL(SD+HD)  |
| Hersteller                       | NordicMubio   |
| Wirt                             | Goat  |
| Kategorie                        | Antikörper  |
| Spezies Reaktivität              | Mouse   |
| Konjugation                      | Unconjugated  |
| Format                           | Antiserum   |
| Spezifität                       | Lambda (Light Chains) - free and bound determinants   |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion   |
| Produktbeschreibung              | The reactivity of the antiserum is directed to the surface and hidden determinants of Ig lambda light chain. In immunoelectrophoresis this antiserum is reacting with polyclonal and monoclonal immunoglobulins of the lambda type, Bence Jones proteins a... |
| Klonalität                       | Monoclonal  |
| Klon-Bezeichnung                 | [Polyclonal]  |

|                        |   |
|------------------------|---|
| Puffer                 | Delipidated, heat inactivated lyophilized whole serum. No preservative added, as it may interfere with the antibody activity It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles.   |
| Quelle                 | A pool of purified Bence Jones lambda proteins isolated from mouse urine. Freund's complete adjuvant is used in the first step of the immunization procedure.   |
| Formel                 | Delipidated, heat inactivated lyophilized whole serum. No preservative added, as it may interfere with the antibody activity  |
| Antibody Type          | Secondary Antibody  |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter center well and 2 $\mu$ l serum samples (neat and serially diluted in 2 mm diameter peripheral wells. When carrying out analyses of Bence Jones proteins or free light chains in urine or serum using an immunodiffusion technique, special attention should be given to their faster rate of diffusion as compared to that of the antibody molecules in the antiserum. If they both start to diffuse into the gel in a double radial diffusion at the same time, an excess of the antigen may easily prevent visible precipitation. To prevent false negative results, the antiserum should be diffuse for about 2 hours before the antigen reservoir is filled. |