

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat anti Rabbit IgG IgA IgM (heavy and light chains), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GAR/IG/TRITC

Artikelname	Goat anti Rabbit IgG IgA IgM (heavy and light chains), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAR/IG/TRITC
Hersteller Artikelnummer	GAR/Ig/TRITC
Alternativnummer	NMB-GAR/IG/TRITC
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Rabbit
Konjugation	TRITC
Format	IgG
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins isotypes It does not react with any non-immunoglobulin in rabbit serum, as tested by immunoelectrophoresis and double radial immunodiffusion. To ident...
Klonalität	Monoclonal

Klon-Bezeichnung	[Polyclonal]
Puffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow
Quelle	Purified immunoglobulin fractions containing IgG, IgA and IgM isolated from pooled rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formulierung	TRITC
Formel	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.