

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat anti Rat Fab, Clone: [Polyclonal], Monoclonal NMB-GARA/FAB

|                                  |   |
|----------------------------------|---|
| Artikelname                      | Goat anti Rat Fab, Clone: [Polyclonal], Monoclonal  |
| Artikelnummer                    | NMB-GARA/FAB  |
| Hersteller Artikelnummer         | GARa/Fab  |
| Alternativnummer                 | NMB-GARA/FAB  |
| Hersteller                       | NordicMubio   |
| Wirt                             | Goat  |
| Kategorie                        | Antikörper  |
| Spezies Reaktivität              | Rat   |
| Konjugation                      | Unconjugated  |
| Format                           | Antiserum   |
| Spezifität                       | IgG (F(ab') <sub>2</sub> )  |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion   |
| Produktbeschreibung              | The antiserum is reacting with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab) <sub>2</sub> subunits and with free light chains of kappa and lambda type as tested in immunoelectrophoresis and dou... |
| Klonalität                       | Monoclonal  |
| Klon-Bezeichnung                 | [Polyclonal]  |

|                        |   |
|------------------------|---|
| Puffer                 | Delipidated, heat inactivated lyophilized stable whole serum No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins added. It is reconstituted by adding 1 ml sterile di   |
| Quelle                 | Purified Fab of normal rat IgG. Freund's complete adjuvant is used in the first step of the immunization procedure.   |
| Formel                 | Delipidated, heat inactivated lyophilized stable whole serum No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins added.   |
| Antibody Type          | Secondary Antibody  |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and diluted) in 2 mm diameter peripheral wells. |