

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Rat IgE (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IGE(FC)/FITC

| | |
|----------------------------------|---|
| Artikelname | Goat anti Rat IgE (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GARA/IGE(FC)/FITC |
| Hersteller Artikelnummer | GARa/IgE(Fc)/FITC |
| Alternativnummer | NMB-GARA/IGE(FC)/FITC |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Spezies Reaktivität | Rat |
| Konjugation | FITC |
| Format | IgG |
| Spezifität | IgE |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc subunit of the IgE molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial in immunodiffusion using various antiserum concentrations against rat serum, a single p... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |

| | |
|------------------------|---|
| Puffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down |
| Quelle | Pools of purified homogenous IgE isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure |
| Formulierung | FITC |
| Formel | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |