

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Rat IgG1 IgG2a IgG2b IgG2c (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GARA/IGG(FC)

Artikelname	Goat anti Rat IgG1 IgG2a IgG2b IgG2c (Fc specific), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GARA/IGG(FC)
Hersteller Artikelnummer	GARa/IgG(Fc)
Alternativnummer	NMB-GARA/IGG(FC)
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Rat
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule including all subclasses. In immunoelectrophoresis and double radial immunodiffusion, using various antiserum concentrations against normal rat plasma and serum, a char...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins adde
Quelle	Purified normal IgG, including all known subclasses, isolated from pooled rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins add
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.