

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Rat IgG2b (subclass specific), Clone: [Polyclonal], Monoclonal NMB-GARA/IGG2B

Artikelname	Goat anti Rat IgG2b (subclass specific), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GARA/IGG2B
Hersteller Artikelnummer	GARa/IgG2b
Alternativnummer	NMB-GARA/IGG2B
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Rat
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG2b (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc portion of the subclass IgG2b. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in at serum, as tested by immunoelectrophoresis and double radia...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protein
Quelle	Pools of purified homogenous IgG2b isolated from pooled rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.