

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Rat IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IGM(FC)/TRITC

Artikelname	Goat anti Rat IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GARA/IGM(FC)/TRITC
Hersteller Artikelnummer	GARa/IgM(Fc)/TRITC
Alternativnummer	NMB-GARA/IGM(FC)/TRITC
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Rat
Konjugation	TRITC
Format	IgG
Spezifität	IgM (μ)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with purified IgG, IgG/Fab fragments and IgA or any non-Ig protein in rat serum, as tested by imm...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow
Quelle	Purified pools of homogenous IgM isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formulierung	TRITC
Formel	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.