

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat anti Swine IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GASW/IGG(H+L)

Artikelname	Goat anti Swine IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GASW/IGG(H+L)
Hersteller Artikelnummer	GASw/IgG(H+L)
Alternativnummer	NMB-GASW/IGG(H+L)
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Porcine
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Feline,Gallus,Canine,Guinea pig,Hamster (all),Equine,Monkey,Mouse,Rabbit,Rat,Sheep
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of both subclasses, IgG1 and IgG2. In immunoelectrophoresis against swine serum, a single characteristic precipitin line is usually observed, representing primarily Igg2. The IgG1...
Klonalität	Monoclonal

Klon-Bezeichnung	[Polyclonal]
Puffer	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Quelle	Purified normal IgG isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and diluted) in 2 mm diameter peripheral wells.