

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Mouse anti CD22, conjugated to PE, IgG1, Clone: [RFB4], Monoclonal NMB-GM-4053

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Mouse anti CD22, conjugated to PE, IgG1, Clone: [RFB4], Monoclonal                                                                                                                                                                                            |
| Artikelnummer            | NMB-GM-4053                                                                                                                                                                                                                                                   |
| Hersteller Artikelnummer | GM-4053                                                                                                                                                                                                                                                       |
| Alternativnummer         | NMB-GM-4053                                                                                                                                                                                                                                                   |
| Hersteller               | NordicMubio                                                                                                                                                                                                                                                   |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF                                                                                                                                                                                                                                                        |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Konjugation              | PE                                                                                                                                                                                                                                                            |
| Produktbeschreibung      | The epitope recognized by antibody RFB4 is a 135 kDa type I integral membrane protein expressed by human B-cells. Precursor B-cells are surface-CD22 negative, but cytoplasmic CD22 positive. Mature B-lymphocytes express CD22 also on their surface. The... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [RFB4]                                                                                                                                                                                                                                                        |
| Isotyp                   | IgG1                                                                                                                                                                                                                                                          |
| UniProt                  | <a href="#">P20273</a>                                                                                                                                                                                                                                        |

|              |                                                                                                              |
|--------------|--------------------------------------------------------------------------------------------------------------|
| Puffer       | 2 ml of PE-conjugated anti CD22 (clone RFB4) in PBS pH 7.2, 1% BSA, and 0.05% NaN3, approximately 100 tests. |
| Reinheit     | Purified by Chromatography                                                                                   |
| Formulierung | PE                                                                                                           |
| Formel       | PBS pH 7.2, 1% BSA, 0.05% NaN3                                                                               |

Anwendungsbeschreibung

Staining Procedure for Surface CD22: Direct Immunofluorescence (Staining Procedure) Nordic-MUBio fluorochrome labeled antibodies are designed for use with either whole blood or isolated mononuclear cell (MNC) preparations Proposed staining procedure for whole blood in short: - For each sample add 50 µl of EDTA anti-coagulated blood to a 3-5 ml tube - Add 20 µl of the appropriate Nordic-MUBio monoclonal antibody conjugate - Incubate the tube for 15 minutes at 4C or at room temperature in the dark - Add 100 µl Nordic-MUBio-LYSE (Cat.No. GAS-003) to each tube and incubate for 10 minutes at room temperature - Add 3-4 ml of distilled water and vortex, incubate for 5-10 minutes at room temperature - Centrifuge tube for 5 minutes at 300 g - Aspirate supernatant and resuspend pellet in 0.3 ml of sheath fluid - Analyze immediately or store samples at 2-8 C in the dark and analyze within 24 hours For "No-Wash protocol please refer to [www.nordicmubio.com](http://www.nordicmubio.com) Proposed staining procedure for MNC in short: - Carefully add 20 µl antibody conjugate and 50-100 µl MNC to the bottom of a tube - Vortex at low speed for 1-2 seconds - Incubate for 15-30 minutes at 2-8C or at room temperature - Centrifuge tubes for 5 minutes at 300 g - Remove supernatant, resuspend cells in 2-5 ml of phosphate buffered saline (PBS) and centrifuge cells again for 5 minutes at 300 g - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1 % formaldehyde and store them at 2-8C in the dark. - Analyze fixed cells within 24 hours

Indirect Immunofluorescence (Staining Procedure) - Mix 20 µl Nordic-MUBio purified antibody with 50 µl whole blood or MNC suspension - Incubate for 15 minutes at 2-8C - Wash cells with phosphate buffered saline (PBS) - Add to cell pellet 20 µl of affinity purified, fluorochrome labeled F(ab)2 anti mouse Ig antibodies - Incubate for 15 minutes at 2-8C - Wash cells with phosphate buffered saline (PBS) or proceed as described for direct staining

Staining Procedure for Cytoplasmatic CD22: Permeabilization and Staining Procedure - In combination with our Permeabilization Kit FIX&PERM (Cat. No. GAS-002) intracellular CD22 can be easily stained in cell suspensions. - For each sample to be analyzed add 50 µl of whole blood, bone marrow or mononuclear cell suspension in a 5ml tube - Add 100 µl of Reagent A (Fixation Medium, stored and used at room temperature) - Incubate for 15 minutes at room temperature - Add 5ml phosphate buffered saline and centrifuge cells for 5 minutes at 300 g - Remove supernatant and add to cell pellet 100 µl Reagent B (Permeabilization Medium) and 20 µl of the CD22 monoclonal antibody conjugate - Vortex at low speed for 1-2 seconds - Incubate for 15 minutes at room temperature - Wash cells with phosphate buffered saline as described above - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1.0 % formaldehyde and store them at 2- 8C in the dark. - Analyze fixed cells within 24 hours.

