

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti CD68 (Macrosialin), conjugated to FITC, IgG1, Clone: [Ki-M7], Monoclonal NMB-GM-4152

Artikelname	Mouse anti CD68 (Macrosialin), conjugated to FITC, IgG1, Clone: [Ki-M7], Monoclonal
Artikelnummer	NMB-GM-4152
Hersteller Artikelnummer	GM-4152
Alternativnummer	NMB-GM-4152
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC
Spezies Reaktivität	Human
Konjugation	FITC
Produktbeschreibung	CD68 (macrosialin) is a type-one membrane protein with significant sequence homology to a family of lysosomal-associated glycoproteins (lamp). CD68 molecules are predominantly located intracellularly in cytoplasmic granules but can also be detected ...
Klonalität	Monoclonal
Klon-Bezeichnung	[Ki-M7]
Isotyp	IgG1
UniProt	P34810

Puffer	Purified by Chromatography, Storage buffer: PBS pH 7.2, 1% BSA, 0.05% NaN3
Reinheit	Purified by Chromatography
Formulierung	FITC
Formel	PBS pH 7.2, 1% BSA, 0.05% NaN3
Anwendungsbeschreibung	<p>Permeabilization and Staining Procedure - In combination with our Permeabilization Kit FIX&PERM? (Cat. No. GAS-002) intracellular CD68 can be easily stained in cell suspensions. - For each sample to be analyzed add 50 µl of whole blood, bone marrow or mononuclear cell suspension in a 5ml tube - Add 100 µl of Reagent A (Fixation Medium, stored and used at room temperature) - Incubate for 15 minutes at room temperature - Add 5 ml phosphate buffered saline and centrifuge cells for 5 minutes at 300 g - Remove supernatant and add to cell pellet 100µl Reagent B (Permeabilization Medium) and 20 µl of the CD68 monoclonal antibody conjugate - Vortex at low speed for 1-2 seconds - Incubate for 15 minutes at room temperature - Wash cells with phosphate buffered saline as described above - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1.0 % formaldehyde and store them at 2-8C in the dark. Analyze fixed cells within 24 hours</p>