

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit anti Bovine erythrocyte, Clone: [Polyclonal], Monoclonal NMB-RAB/ERY/7S

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit anti Bovine erythrocyte, Clone: [Polyclonal], Monoclonal   |
| Artikelnummer            | NMB-RAB/ERY/7S  |
| Hersteller Artikelnummer | RAB/Ery/7S  |
| Alternativnummer         | NMB-RAB/ERY/7S  |
| Hersteller               | NordicMubio   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Produktbeschreibung      | To sensitize cells in haemolytic complement assays and for screening of T and T receptor-bearing lymphocytes....  |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [Polyclonal]  |
| Puffer                   | Purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. IgG protein concentration 10 mg/ml. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 ml sterile dis |
| Quelle                   | Washed whole erythrocytes were prepared from fresh blood and used for intravenous immunization. Freund's complete adjuvant is used in the first step of the immunization procedure  |
| Formel                   | Purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. IgG protein concentration 10 mg/ml. No foreign proteins added.   |

|                        |  |
|------------------------|--|
| Anwendungsbeschreibung | Indirect immunofluorescence,ELISA,Dot blot,Immunoblotting.<br>Haemolytic titre, at which 50% lysis occurs following mixing of equal volumes of antiserum, guinea pig complement (at a 1:60 dilution) and 2% RBC, not less than 1:500 when tested with bovine erythrocytes. |
|------------------------|--|