

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Cat IgG (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RACA/IGG(FC)

| | |
|----------------------------------|---|
| Artikelname | Rabbit anti Cat IgG (Fc specific), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-RACA/IGG(FC) |
| Hersteller Artikelnummer | RACa/IgG(Fc) |
| Alternativnummer | NMB-RACA/IGG(FC) |
| Hersteller | NordicMubio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Spezies Reaktivität | Feline |
| Konjugation | Unconjugated |
| Format | Antiserum |
| Spezifität | IgG (Fc) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal cat serum a single precipitin line is obtained which shows a ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |

| | |
|------------------------|--|
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized |
| Quelle | Highly purified normal IgG isolated from pooled cat serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. |
| Antibody Type | Secondary Antibody |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. |