

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Duck IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RADU/IGG(H+L)

Artikelname	Rabbit anti Duck IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RADU/IGG(H+L)
Hersteller Artikelnummer	RADu/IgG(H+L)
Alternativnummer	NMB-RADU/IGG(H+L)
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IP
Spezies Reaktivität	Duck
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Feline,Gallus,Canine,Goat,Equine,Human,Monkey,Mouse,Rat,S heep,Porcine
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in duck serum, as ...

Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added. It is reconstituted by adding 1 ml s
Quelle	Purified normal IgG isolated from pooled duck serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells. The optimum working dilution of the antibody for any specific application should be established by titration.