

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Human heparin cofactor II, Clone: [Polyclonal], Monoclonal NMB-RAHU/HCFII

Artikelname	Rabbit anti Human heparin cofactor II, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RAHU/HCFII
Hersteller Artikelnummer	RAHu/HCFII
Alternativnummer	NMB-RAHU/HCFII
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Produktbeschreibung	The reactivity of the antiserum is restricted to heparin cofactor II. In bidimensional electrophoresis against fresh normal human plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with pu...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
UniProt	P05546
Puffer	Delidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized a

Quelle	<p>Heparin cofactor II is a glycoprotein synthesized in the liver and has a molecular weight of 65,000. It is a plasma serine inhibitor of thrombin. Antithrombin III is more slowly inhibited. A hereditary form of inactive heparin cofactor II has been described</p>
Formel	<p>Delidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.</p>
Anwendungsbeschreibung	<p>Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl in a 3 mm diameter center well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. For use in electroimmunodiffusion use a concentration of 2% antiserum in the gel. Measured by quantitative precipitin analysis. The amount of heparin cofactor II precipitated by 1 ml antiserum is approximately 90 µg/ml.</p>