

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Human IgG3 (subclass specific), Clone: [Polyclonal], Monoclonal NMB-RAHU/IGG3

Artikelname	Rabbit anti Human IgG3 (subclass specific), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RAHU/IGG3
Hersteller Artikelnummer	RAHu/IgG3
Alternativnummer	NMB-RAHU/IGG3
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Spezies Reaktivität	Human
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG3 (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is restricted to subclass specific determinants on IgG3. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal human serum a single precipitin line is obtained which...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized
Quelle	Highly purified homogenous IgG3 isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In double radial immunodiffusion (Ouchterlony) serum or plasma samples are prepared in serial two-fold dilutions from 1:8 to 1: 128 and placed in the peripheral wells of the rosette arrangement. One well is filled with pooled human serum or another control. The center wells are filled with the subclass specific antiserum and the sera are allowed to diffuse at ambient temperature for 16-24 hours. Frequently the precipitin lines become visible only after staining.