

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Human IgG IgA IgM IgD IgE (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAHU/IG(GAMDE)

| | |
|----------------------------------|---|
| Artikelname | Rabbit anti Human IgG IgA IgM IgD IgE (heavy and light chains), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-RAHU/IG(GAMDE) |
| Hersteller Artikelnummer | RAHu/Ig(GAMDE) |
| Alternativnummer | NMB-RAHU/IG(GAMDE) |
| Hersteller | NordicMubio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Konjugation | Unconjugated |
| Format | Antiserum |
| Spezifität | IgG+IgM+IgA+IgD+IgE (H+L) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | The reactivity of the antiserum is directed to the heavy and light chains of the immunoglobulin isotypes. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunoelectro... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |

| | |
|------------------------|---|
| Puffer | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s |
| Quelle | Purified normal Ig fractions isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit |
| Antibody Type | Secondary Antibody |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells. |