

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Human platelet factor 4, Clone: [Polyclonal], Monoclonal NMB-RAHU/PF4

Artikelname	Rabbit anti Human platelet factor 4, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RAHU/PF4
Hersteller Artikelnummer	RAHu/PF4
Alternativnummer	NMB-RAHU/PF4
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Produktbeschreibung	The reactivity of the antiserum is restricted to platelet factor 4 as tested in immunoelectrophoresis and radial immunodiffusion. A characteristic single precipitin line is obtained with normal serum which shows a reaction of identity with the precip...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
UniProt	P02776
Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum Sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized antiserum
Quelle	Platelet factor 4 is a low molecular weight (MW 7,800) heat stable protein, secreted by alpha granules of the platelets. Purified platelet factor 4 isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization p

Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum Sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In electroimmunodiffusion the amount of antiserum required in the agarose gel is usually between 1 and 2%, depending on the test arrangement. The amount of platelet factor 4 precipitated by 1 ml antiserum is approximately 0.1 mg.