

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Sheep IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RASH/IGM(FC)

Artikelname	Rabbit anti Sheep IgM (Fc specific), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RASH/IGM(FC)
Hersteller Artikelnummer	RASH/IgM(Fc)
Alternativnummer	NMB-RASH/IGM(FC)
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Spezies Reaktivität	Sheep
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgM (μ)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal sheep serum a single precipitin line is obtained which shows ...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized
Quelle	Highly purified normal IgM isolated from pooled sheep serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 μ l serum, plasma or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single radial immunodiffusion and electroimmunodiffusion use 0.5 to 1.0 percent antiserum in the agar gel.