

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Sheep IgM (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-RASH/IGM(FC)/FITC

| | |
|----------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Rabbit anti Sheep IgM (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-RASH/IGM(FC)/FITC |
| Hersteller Artikelnummer | RASH/IgM(Fc)/FITC |
| Alternativnummer | NMB-RASH/IGM(FC)/FITC |
| Hersteller | NordicMubio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Spezies Reaktivität | Sheep |
| Konjugation | FITC |
| Format | IgG |
| Spezifität | IgM (μ) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in sheep serum, as tested by immunoelectrophoresis and double radial immu... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer | Fluorescein-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, |
| Quelle | Purified normal IgM isolated from pooled sheep serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formulierung | FITC |
| Formel | Fluorescein-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |