

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Human RANTES Antibody, Mouse, Monoclonal ABT-ABG10473-U500

Artikelname	Anti-Human RANTES Antibody, Mouse, Monoclonal
Artikelnummer	ABT-ABG10473-U500
Hersteller Artikelnummer	ABG10473-U500
Alternativnummer	ABT-ABG10473-U500-500UG
Hersteller	Abcepta
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Monoclonal
Reinheit	Produced in BALB/c mice using highly pure (>98%) human RANTES as the immunizing antigen. This IgG2bK antibody was purified from ascites fluid by Protein A affinity chromatography.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS.
Antibody Type	Monoclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect hRANTES by Western Blot analysis this antibody can be used at a concentration of 0.5-1.0 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hRANTES is 0.25-2.0 ng/lane, under reducing conditions and 0.25-0.50 ng/lane, under non-reducing conditions.. Sandwich: In a sandwich ELISA (assuming 100µl/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 500 pg/ml of recombinant human RANTES when used with BioGems biotinylated antigen affinity purified anti-human RANTES (60-285BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human tonsil. The recommended concentration is 10.0 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 citrate buffer is recommended. Optimal concentrations and conditions may vary. Protocol and staining provided by Dr. Lauren Binge, Laboratory of Prof. Charles Mackay, Monash Univeristy, Australia.

 *Additional Immunostaining data available. Please contact Tech Support for information.. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Human RANTES (100 ng/ml), a concentration of 3.0-5.0 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.