

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human SDF-1alpha Antibody, Rabbit, Polyclonal ABT-ABG10500-U050

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Human SDF-1alpha Antibody, Rabbit, Polyclonal  |
| Artikelnummer            | ABT-ABG10500-U050   |
| Hersteller Artikelnummer | ABG10500-U050   |
| Alternativnummer         | ABT-ABG10500-U050-50UG  |
| Hersteller               | Abcepta   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, IHC, WB  |
| Spezies Reaktivität      | Human   |
| Klonalität               | Polyclonal  |
| Reinheit                 | Produced from sera of rabbits pre-immunized with highly pure recombinant Human SDF-1alpha. Anti-Human SDF-1alpha specific antibody was purified by affinity chromatography employing immobilized Human SDF-1alpha matrix. |
| Formulierung             | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.  |
| Antibody Type            | Polyclonal Antibody   |

Anwendungsbeschreibung

WesternBlot: To detect Human SDF-1alpha by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human SDF-1alpha is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions..

Sandwich: To detect Human SDF-1alpha by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human SDF-1alpha (60-294BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human SDF-1alpha..

Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human colorectal adenocarcinoma. The recommended concentration is 0.5 µg/ml with an overnight incubation at 4C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary. <br /> <br /> <span style=font-size: 10px,>\*Additional Immunostaining data available. Please contact Tech Support for information. . Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of Human SDF-1alpha (100 ng/ml), a concentration of 2-4 µg/ml of this antibody is required..

Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.