

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Biotinylated Anti-Human SDF-1alpha Antibody, Rabbit, Polyclonal ABT-ABG10502-U050

| | |
|--------------------------|---|
| Artikelname | Biotinylated Anti-Human SDF-1alpha Antibody, Rabbit, Polyclonal |
| Artikelnummer | ABT-ABG10502-U050 |
| Hersteller Artikelnummer | ABG10502-U050 |
| Alternativnummer | ABT-ABG10502-U050-50UG |
| Hersteller | Abcepta |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA, WB |
| Spezies Reaktivität | Human |
| Klonalität | Polyclonal |
| Reinheit | Produced from sera of rabbits pre-immunized with highly pure recombinant Human SDF-1alpha. Anti-Human SDF-1alpha specific antibody was purified by affinity chromatography and then biotinylated. |
| Formulierung | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2. |
| Antibody Type | Polyclonal Antibody |

| | |
|------------------------|--|
| Anwendungsbeschreibung | <p>WesternBlot: To detect Human SDF-1alpha by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human SDF-1alpha is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions..</p> <p>Sandwich: To detect Human SDF-1alpha by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems Polyclonal Anti-Human SDF-1alpha (60-294P) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human SDF-1alpha..</p> <p>Direct: To detect Human SDF-1alpha by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human SDF-1alpha..</p> <p>Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.</p> |
|------------------------|--|