

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Murine sRANK Ligand Antibody, Rabbit, Polyclonal ABT-ABG10515-U050

Artikelname	Anti-Murine sRANK Ligand Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10515-U050
Hersteller Artikelnummer	ABG10515-U050
Alternativnummer	ABT-ABG10515-U050-50UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, WB
Spezies Reaktivität	Mouse
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant msRANKL. Anti-Murine sRANKL specific antibody was purified by affinity chromatography employing immobilized msRANKL matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect msRANKL by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant msRANKL is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect msRANKL by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine sRANKL (61-113BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant msRANKL.. Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of msRANKL (50.0 ng/ml), a concentration of 0.05 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.