

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Human beta-NGF Antibody, Rabbit, Polyclonal ABT-ABG10522-U050

| | |
|--------------------------|---|
| Artikelname | Anti-Human beta-NGF Antibody, Rabbit, Polyclonal |
| Artikelnummer | ABT-ABG10522-U050 |
| Hersteller Artikelnummer | ABG10522-U050 |
| Alternativnummer | ABT-ABG10522-U050-50UG |
| Hersteller | Abcepta |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC, WB |
| Spezies Reaktivität | Human |
| Klonalität | Polyclonal |
| Reinheit | Produced from sera of rabbits pre-immunized with highly pure recombinant Human beta-NGF. Anti-Human beta-NGF specific antibody was purified by affinity chromatography employing immobilized Human beta-NGF matrix. |
| Formulierung | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2. |
| Antibody Type | Polyclonal Antibody |

Anwendungsbeschreibung

WesternBlot: To detect Human beta-NGF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human beta-NGF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect Human beta-NGF by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human beta-NGF (60-247BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human beta-NGF.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human pancreas adenocarcinoma. The recommended concentration is 0.5 mg/ml with an overnight incubation at 4C. An HRP-labeled polymer detection system was used with a DAB chromogen. Optimal results were achieved without antigen retrieval. Optimal concentrations and conditions may vary.. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Human beta-NGF (1 ng/ml), a concentration of 0.05-0.08 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.