

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### HDAC6 Antibody, IgG2a, Clone: [PCRP-HDAC6-1A4], Mouse, Monoclonal NBT-10013-MSM1-P0

|                          |   |
|--------------------------|---|
| Artikelname              | HDAC6 Antibody, IgG2a, Clone: [PCRP-HDAC6-1A4], Mouse, Monoclonal   |
| Artikelnummer            | NBT-10013-MSM1-P0   |
| Hersteller Artikelnummer | 10013-MSM1-P0   |
| Alternativnummer         | NBT-10013-MSM1-P0-20,NBT-10013-MSM1-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human HDAC6 protein   |
| Produktbeschreibung      | In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought a... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PCRP-HDAC6-1A4]  |
| Molekulargewicht         | 160kDa  |
| Isotyp                   | IgG2a   |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">10013</a>  |
| UniProt                | <a href="#">Q9UBN7</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.                               |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunoprecipitation (1-2ug per 100-500ug of total protein), ,Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), ,Optimal dilution for a specific application should be determined. |