

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant p21WAF1 (Tumor Suppressor Protein) Antibody, Clone: [CIP1/2275R], Rabbit, Monoclonal NBT-1026-RBM7-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant p21WAF1 (Tumor Suppressor Protein) Antibody, Clone: [CIP1/2275R], Rabbit, Monoclonal |
| Artikelnummer | NBT-1026-RBM7-P0 |
| Hersteller Artikelnummer | 1026-RBM7-P0 |
| Alternativnummer | NBT-1026-RBM7-P0-20,NBT-1026-RBM7-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CDKN1A protein |
| Produktbeschreibung | This MAb recognizes a 21kDa protein, identified as the p21WAF1 tumor suppressor protein. This MAb is highly specific to p21 and shows no cross-reaction with other closely related mitotic inhibitors. p21WAF1 is a specific inhibitor of cdks and a tumor... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CIP1/2275R] |
| Molekulargewicht | 21kDa |
| NCBI | 1026 |

| | |
|------------------------|---|
| UniProt | P38936 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 min |