

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Clusterin / Apolipoprotein J (APO-J) Antibody, IgG2c, Clone: [CLU/4727], Mouse, Monoclonal NBT-1191-MSM7-P1ABX

| | |
|--------------------------|---|
| Artikelname | Clusterin / Apolipoprotein J (APO-J) Antibody, IgG2c, Clone: [CLU/4727], Mouse, Monoclonal |
| Artikelnummer | NBT-1191-MSM7-P1ABX |
| Hersteller Artikelnummer | 1191-MSM7-P1ABX |
| Alternativnummer | NBT-1191-MSM7-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa150-300) of human CLU protein (exact sequence is proprietary) |
| Produktbeschreibung | Clusterin, also designated complement lysis inhibitor (CLI), apolipoprotein J (APOJ), sulfated glycoprotein 2 (SGP2), SP40 and testosterone-repressed prostate message 2 (TRPM2), is a secretory, heterodimeric glycoprotein that influences immune regula... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CLU/4727] |
| Molekulargewicht | 70kDa (precursor), 36-39kDa (v(TM)), 34-36kDa (v°a) |

| | |
|------------------------|---|
| Isotyp | IgG2c |
| NCBI | 1191 |
| UniProt | P10909 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (2-4ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for |