

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Catenin, beta (p120) Antibody, IgG1, Clone: [CTNNB1/1509], Mouse, Monoclonal NBT-1499-MSM9-P1ABX

| | |
|--------------------------|---|
| Artikelname | Catenin, beta (p120) Antibody, IgG1, Clone: [CTNNB1/1509], Mouse, Monoclonal |
| Artikelnummer | NBT-1499-MSM9-P1ABX |
| Hersteller Artikelnummer | 1499-MSM9-P1ABX |
| Alternativnummer | NBT-1499-MSM9-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Recombinant human beta-Catenin (p120) protein fragment |
| Produktbeschreibung | Beta-catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. In normal tissues, beta-catenin is localized to the membrane of epithelial cells, consistent with its role ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CTNNB1/1509] |
| Molekulargewicht | 92kDa |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| NCBI | 1499 |
| UniProt | P35222 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution fo |