

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cathepsin K (Marker of Tumor Invasiveness) Antibody, IgG1, Clone: [CTSK/2792], Mouse, Monoclonal NBT-1513-MSM2-P1ABX

| | |
|--------------------------|---|
| Artikelname | Cathepsin K (Marker of Tumor Invasiveness) Antibody, IgG1, Clone: [CTSK/2792], Mouse, Monoclonal |
| Artikelnummer | NBT-1513-MSM2-P1ABX |
| Hersteller Artikelnummer | 1513-MSM2-P1ABX |
| Alternativnummer | NBT-1513-MSM2-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human Cathepsin K protein (around aa 163-274) (exact sequence is proprietary) |
| Produktbeschreibung | The protein encoded by this gene is a lysosomal cysteine proteinase involved in bone remodeling and resorption. This protein, which is a member of the peptidase C1 protein family, is predominantly expressed in osteoclasts. However, the encoded protei... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CTSK/2792] |
| Molekulargewicht | 39kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 1513 |
| UniProt | P43235 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution |