

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TdT / DNA Nucleotidylexotransferase (Acute Lymphoblastic Leukemia Marker) Antibody, IgG1, Clone: [TDT/1393], Mouse, Monoclonal NBT-1791-MSM3-P0

| | |
|--------------------------|---|
| Artikelname | TdT / DNA Nucleotidylexotransferase (Acute Lymphoblastic Leukemia Marker) Antibody, IgG1, Clone: [TDT/1393], Mouse, Monoclonal |
| Artikelnummer | NBT-1791-MSM3-P0 |
| Hersteller Artikelnummer | 1791-MSM3-P0 |
| Alternativnummer | NBT-1791-MSM3-P0-20,NBT-1791-MSM3-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 52-192) of human DNTP protein (exact sequence is proprietary) |
| Produktbeschreibung | Terminal deoxynucleotidyl transferase (TdT) is an unusual deoxynucleotide polymerizing enzyme with a molecular weight of about 58 kDa found normally only in B- and T-cell lymphoblasts/prelymphocytes. TdT generates antigen receptor diversity by synthe... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [TDT/1393] |
| Molekulargewicht | 58kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 1791 |
| UniProt | P04053 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 9 |