

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AIF1 / Iba1 (Microglia Marker) Antibody, IgG1, Clone: [AIF1/1909], Mouse, Monoclonal NBT-199-MSM1-P0

| | |
|--------------------------|---|
| Artikelname | AIF1 / Iba1 (Microglia Marker) Antibody, IgG1, Clone: [AIF1/1909], Mouse, Monoclonal |
| Artikelnummer | NBT-199-MSM1-P0 |
| Hersteller Artikelnummer | 199-MSM1-P0 |
| Alternativnummer | NBT-199-MSM1-P0-20,NBT-199-MSM1-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Purified fragment of human recombinant AIF1 protein (around aa 1-146) (exact sequence is proprietary) |
| Produktbeschreibung | AIF1 is a cytoplasmic, calcium-binding protein that is thought to play a role in macrophage activation and function. AIF1, containing two EF domains, is induced by cytokines and Interferons. In an unstimulated state, AIF1 colocalizes with actin, and ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [AIF1/1909] |
| Molekulargewicht | 17kDa |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| NCBI | 199 |
| UniProt | P55008 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |