

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AKT1 (Prognostic Marker for Neuroendocrine Tumors) Antibody, IgG2b, Clone: [AKT1/2552], Mouse, Monoclonal NBT-207-MSM2-P0

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | AKT1 (Prognostic Marker for Neuroendocrine Tumors) Antibody, IgG2b, Clone: [AKT1/2552], Mouse, Monoclonal |
| Artikelnummer | NBT-207-MSM2-P0 |
| Hersteller Artikelnummer | 207-MSM2-P0 |
| Alternativnummer | NBT-207-MSM2-P0-20,NBT-207-MSM2-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human AKT1 protein (around aa 85-189) (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of 62kDa, which is identified as AKT1. The serine/threonine kinase Akt family contains several members, including Akt1 (also designated PKB or RacPK), Akt2 (also designated PKB tyrosine residues 740 and 751, which bind the subuni... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [AKT1/2552] |
| Molekulargewicht | 62kDa |

| | |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG2b |
| NCBI | 207 |
| UniProt | P31749 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml),Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes) |