

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Alkaline Phosphatase (Placental) / PLAP (Germ Cell Tumor Marker) Antibody, IgG2b, Clone: [PL8-F6], Mouse, Monoclonal NBT-250-MSM5-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Alkaline Phosphatase (Placental) / PLAP (Germ Cell Tumor Marker) Antibody, IgG2b, Clone: [PL8-F6], Mouse, Monoclonal  |
| Artikelnummer            | NBT-250-MSM5-P0   |
| Hersteller Artikelnummer | 250-MSM5-P0   |
| Alternativnummer         | NBT-250-MSM5-P0-20,NBT-250-MSM5-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Purified human PLAP protein   |
| Produktbeschreibung      | Reacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PL8-F6]  |
| Molekulargewicht         | 70kDa   |
| Isotyp                   | IgG2b   |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">250</a>   |
| UniProt                | <a href="#">P05187</a>  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Frozen & Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT)(No special pretreatment is required for the immunohistochemical staining of |