

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant PD-L1 / PDCD1LG1 / CD274 / B7-H1 (Cancer Immunotherapy Target) Antibody, Clone: [PDL1/4280R], Rabbit, Monoclonal NBT-29126-RBM10-P1ABX

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant PD-L1 / PDCD1LG1 / CD274 / B7-H1 (Cancer Immunotherapy Target) Antibody, Clone: [PDL1/4280R], Rabbit, Monoclonal  |
| Artikelnummer            | NBT-29126-RBM10-P1ABX   |
| Hersteller Artikelnummer | 29126-RBM10-P1ABX   |
| Alternativnummer         | NBT-29126-RBM10-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | KLH-conjugated linear peptide corresponding to human PD-L1  |
| Produktbeschreibung      | PD-L1 is a checkpoint regulator in immune cells, it is expressed on immune or non-hematopoietic cells. Expression of the protein is seen during pregnancy where it has a role in suppressing the immune system. PD-L1 induces an inhibitory signal in acti... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PDL1/4280R]  |
| Molekulargewicht         | 37-50kDa  |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">29126</a>  |
| UniProt                | <a href="#">Q9NZQ7</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Recombinant Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |