

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Alpha-2-Macroglobulin Antibody, IgG1, Clone: [A2M/3623], Mouse, Monoclonal NBT-2-MSM3-P1ABX

| | |
|--------------------------|---|
| Artikelname | Alpha-2-Macroglobulin Antibody, IgG1, Clone: [A2M/3623], Mouse, Monoclonal |
| Artikelnummer | NBT-2-MSM3-P1ABX |
| Hersteller Artikelnummer | 2-MSM3-P1ABX |
| Alternativnummer | NBT-2-MSM3-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human recombinant A2M protein fragment (around aa 604-748) (exact sequence is proprietary) |
| Produktbeschreibung | (TM)-2-Macroglobulin ((TM)-2M) is a homotetrameric serum protein consisting of four identical subunits that form dimers through disulfide bonds. Initially, (TM)-2M was characterized as a pan-proteinase inhibitor that was able to bait proteinases into... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [A2M/3623] |
| Molekulargewicht | 185kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 2 |
| UniProt | P01023 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |