

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant IgM (Immunoglobulin Mu Heavy Chain) (B-Cell Marker) Antibody, IgG1, Clone: [rIGHM/2558], Mouse, Monoclonal NBT-3507-MSM9-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant IgM (Immunoglobulin Mu Heavy Chain) (B-Cell Marker) Antibody, IgG1, Clone: [rIGHM/2558], Mouse, Monoclonal  |
| Artikelnummer            | NBT-3507-MSM9-P1ABX   |
| Hersteller Artikelnummer | 3507-MSM9-P1ABX   |
| Alternativnummer         | NBT-3507-MSM9-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Heavy chain of human IgM  |
| Produktbeschreibung      | Recognizes a protein of 75kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. Monomeric IgM... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rIGHM/2558]  |
| Molekulargewicht         | 50-75kDa  |
| Isotyp                   | IgG1  |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">3507</a>  |
| UniProt                | <a href="#">P01871</a>  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min |