

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Arginase1 (Hepatocellular Carcinoma Marker) Antibody, IgG3, Clone: [ARG1/1125], Mouse, Monoclonal NBT-383-MSM1-P1ABX

| | |
|--------------------------|---|
| Artikelname | Arginase1 (Hepatocellular Carcinoma Marker) Antibody, IgG3, Clone: [ARG1/1125], Mouse, Monoclonal |
| Artikelnummer | NBT-383-MSM1-P1ABX |
| Hersteller Artikelnummer | 383-MSM1-P1ABX |
| Alternativnummer | NBT-383-MSM1-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human ARG1 protein fragment (around aa11-97) (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of 35-38kDa, which is identified as Arginase 1 (ARG1). Arginase is a manganese metallo-enzyme that catalyzes the hydrolysis of arginine to generate ornithine and urea. Arginase I and II are isoenzymes which differ in subcellular ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [ARG1/1125] |
| Molekulargewicht | 35-38kDa |

| | |
|------------------------|---|
| Isotyp | IgG3 |
| NCBI | 383 |
| UniProt | P05089 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blotting (1-2ug/ml), Immunohistology (Formalin-fixed) (2-4ug/ml for 30 minutes at RT) ,(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes), Op |