

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 7 (Glandular and Transitional Epithelial Marker) Antibody, IgG1, Clone: [SPM270], Mouse, Monoclonal NBT-3855-MSM1X-P0

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 7 (Glandular and Transitional Epithelial Marker) Antibody, IgG1, Clone: [SPM270], Mouse, Monoclonal |
| Artikelnummer | NBT-3855-MSM1X-P0 |
| Hersteller Artikelnummer | 3855-MSM1X-P0 |
| Alternativnummer | NBT-3855-MSM1X-P0-20,NBT-3855-MSM1X-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | OTN 11, ovarian carcinoma cell line |
| Produktbeschreibung | It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in mos... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM270] |
| Molekulargewicht | 55kDa |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| NCBI | 3855 |
| UniProt | P08729 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM |