

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Cytokeratin 8 (KRT8) Antibody, Clone: [KRT8/4067R], Rabbit, Monoclonal NBT-3856-RBM17-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant Cytokeratin 8 (KRT8) Antibody, Clone: [KRT8/4067R], Rabbit, Monoclonal |
| Artikelnummer | NBT-3856-RBM17-P0 |
| Hersteller Artikelnummer | 3856-RBM17-P0 |
| Alternativnummer | NBT-3856-RBM17-P0-20,NBT-3856-RBM17-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Keratin preparation from a human carcinoma |
| Produktbeschreibung | Epitope of this MAb is located between aa343-357. Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squam... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT8/4067R] |
| Molekulargewicht | 52.5kDa |
| NCBI | 3856 |

| | |
|------------------------|---|
| UniProt | P05787 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (2-4ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1m |